



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTIONS OF PROBLEMS IN NUMBER THREE.

SOLUTIONS of problems in No. 3 have been received as follows:

From Prof. W. P. Casey, 437, 438; Prof. H. T. Eddy, 440; Prof. A. Hall, 440; E. H. Moore, Jr., 438; Ernest G. Merritt, 438.

437. *By Prof. Casey.*—"AE, AK are two indefinite given straight lines, C and H given points in them, and P a given point in their plane. Req'd to draw through P two straight lines, PB, PD, intersecting AE and AK in R and S, respectively, and containing a given angle RPS, so that $RC \times SH$ may be equal to a given magnitude."


SOLUTION BY THE PROPOSER.

Let AE, AK be two indefinite straight lines given in position; C, H, given points in them, and P a giv'n point in their plane; and let PR, PS be drawn meeting them in R, S, so that $RC \times HS$ may be = a given space, A^2 ; and the $\angle RPS$ = a given angle.

Analysis.—Join PC; it is given in position. Through P, H, draw PO, DHO, parallel to AE, PC, respectively; \therefore O is a given point. Produce RP to meet DO in N. Now $RC \times ON = PC \times CO$ and is therefore given, and $RC \times HS$ is given; therefore the ratio of ON to HS is given. Angle RPS is given, $\therefore \angle NPS$ is given.

Make $\angle OPG = \angle NPS$, then PG is in position and G is a given point. Make $\angle PGF = \angle PON$ = a given angle, therefore GF is in position, and triangles PON and PGF are similar, and $ON : GF :: PO : PG$, i. e., in a given ratio, and the ratio of $ON : HS$ is given, \therefore the ratio of $GF : HS$ is given.

Through H draw HQ parallel to GF, \therefore HQ is in position. Draw SY parallel to HQ, and GY parallel to PS, meeting SY in Y. Through Y draw XYM parallel to AK, and through P draw PQ parallel to AK, \therefore PQ is in position, and Q is a given point. Join HY and produce it to meet PQ in Z and draw HL parallel to PS. Then $GF = SY = HX$, and




$HS = XY$; and as the ratio of $GF : HS$ is given, \therefore the ratio of HX to XY is given; hence $HQ : QZ$ is given and HQ is given, $\therefore QZ$ is given and Q is a given point, $\therefore Z$ is a given point and PZ is a given line. But $QZ \times YL = XY \times ZV$, and $YL = GH$, a given line, also $XY = HS = VP$, $\therefore ZV \times VP = QZ \times GH$, a given space, and ZP is given, $\therefore V$ is a given point and VH is in position, hence PS is in position and so is PR , and the problem is solved geometrically.

438. By Prof. F. H. Loud.—“Two equiangular parallelograms, $OACB$ and $OA'C'B'$ are so placed that the equal angles AOB and $A'O'B'$ coincide. The sides of the former figure are constant, those of the latter are variable, subject to the condition $A'O + OB' : AO + OB :: \text{area } OA'C'B' : \text{area } OACB$. $A'C'$ meets BC in D , and $B'C'$ meets AC in E . Show that DE passes through a fixed point, and determine the point.”

SOLUTION BY ERNEST G. MERRITT, CORNELL UNIV., ITHACA, N. Y.

Put $OB = a$
 $OA = b$, coordinates of the point C , and let the coordinates of C' be $OB' = x'$ and $OA' = y'$.

Then, from the conditions of the problem,



$$x' + y' : a + b = x'y' \sin AOB : ab \sin AOB;$$

$$\therefore (a+b)x'y' = ab(x' + y'), \quad (1)$$

whence

$$y' = \frac{abx'}{(a+b)x' - ab}. \quad (2)$$

The coordinates of D are (a, y') , and of E , (x', b) . Substituting in the equation of a line through two points (x_1, y_1) and (x_2, y_2)

$$\left[y - y_1 = (x - x_1) \frac{y_2 - y_1}{x_2 - x_1} \right],$$

we obtain $y - b = \left(\frac{x - x'}{a - x'} \right) \left(\frac{abx'}{(a+b)x' - ab} - b \right)$, which may be written

$$y - b = (x - x') \frac{\frac{ab}{a+b} - b}{\frac{ab}{a+b} - x'}. \quad (3)$$

But $u = x^2 + m^2y^2$, $v = x^2 - n^2y^2$; therefore

$$F = \pm \int_{a'^2}^{a^2} \int_{\beta'^2}^{\beta^2} \frac{xy du dv}{4xy(m^2+n^2)} = \frac{(a^2-a'^2)(\beta^2-\beta'^2)}{4(m^2+n^2)}.$$

SOLUTION BY PROF. ASAPH HALL.

The ratio of the axes of the curves being given, we may write the eq'ns

$$x^2 + m^2y^2 = u; \quad x^2 - n^2y^2 = v.$$

The product of inertia is given by the integral $\int xy dx dy$, and we have to transform this to the variables u and v . The partial derivatives are,

$$\begin{aligned} \frac{dx}{du} &= \frac{n^2}{2x(m^2+n^2)}; & \frac{dx}{dv} &= \frac{m^2}{2x(m^2+n^2)}; \\ \frac{dy}{du} &= \frac{1}{2y(m^2+n^2)}; & \frac{dy}{dv} &= \frac{-1}{2y(m^2+n^2)}. \end{aligned}$$

Forming the known determinant for the transformation we have,

$$\int xy dx dy = \frac{1}{4(m^2+n^2)} \int du dv = \frac{(a^2-a'^2)(\beta^2+\beta'^2)}{4(m^2+n^2)};$$

since the limits of u are a^2 and a'^2 , and of v , β^2 and β'^2 .

REMARKS ON "NEW RULE FOR CUBE ROOT."—We published on page 98, No. 3, what purports to be a new Rule for Cube Root, and were not aware, at the time, that substantially the same rule had been published before; and we have no doubt the author believed it to be new, and original with him.

The same Rule, in effect, may be found at page 32, Vol. I of the *Mathematical Monthly*, published in Nov., 1859, at Cambridge, Mass. The editor (J. D. Runkle) there says, "In the *Nouvelles Mathématiques* for January, 1858, we find the following method for extracting the cube root of numbers, which ought, on account of its easy application, to be generally used. The editor remarks, in the April number, that the method had previously been given in a work entitled *Calcul pratiques*, in which it is claimed as new. The reader will find the same process, entitled a new method, in the American edition of Young's *Algebra*, published as long ago as 1832. It may also be found in some of our arithmetics; and many teachers undoubtedly already know and use it."

PROBLEMS.

441. By *Wm. Hoover, A. M., Dayton, Ohio.*—A cone revolves around its axis with a known angular velocity. The altitude begins to diminish